

Karnofsky score, HEM 13892

Skattning av aktivitetsnivå

Karnofsky score för vuxna och ungdomar ≥ 16 år:

100 %	normal activity, no complaints, no signs of disease
90 %	capable of normal activity, few symptoms or signs of disease
80 %	normal activity with some difficulty, some symptoms or signs
70 %	caring for self, not capable of normal activity or work
60 %	requiring some help, can take care of most personal requirements
50 %	requires help often, requires frequent medical care
40 %	disabled, requires special care and help
30 %	severely disabled, hospital admission indicated but no risk of death
20 %	very ill, urgently requiring admission, requires supportive measures or treatment
10 %	moribund, rapidly progressive fatal disease processes
0 %	death

Referenser

Karnofsky, DA., Burchenal, JH. (1949). The Clinical Evaluation of Chemotherapeutic Agents in Cancer. *MacLeod CM (Ed), Evaluation of Chemotherapeutic Agents*. Columbia Univ Press. p 196.

Schag, CC., Heinrich, RL., Ganz, PA. Karnofsky performance status revisited: Reliability, validity, and guidelines. *J Clin Oncology* 1984; 2:187-193.

O'Toole, DM., Golden, AM. Evaluating cancer patients for rehabilitation potential. *West J Med*. 1991; 155:384-387.

Oxford Textbook of Palliative Medicine. (1993). Oxford University Press. 1993;109